

Completeness of the $SO(4)$ extended
Bethe Ansatz
for the One-dimensional Hubbard Model

Fabian H.L. Eßler
Vladimir E. Korepin
and
Kareljan Schoutens

*Institute for Theoretical Physics
State University of New York at Stony Brook
Stony Brook, NY 11794-3840, U.S.A.*

Abstract

We show how to construct a complete set of eigenstates of the hamiltonian of the one-dimensional Hubbard model on a lattice of even length L . This is done by using the nested Bethe Ansatz *and* the $SO(4)$ symmetry of the model. We discuss in detail how the counting of independent eigenstates is carried out.

1 Introduction

An important ingredient in the search for a theory of high- T_c superconductors is the analysis of strongly correlated electron systems ^[1]. A prototype model for these is the Hubbard model. Especially interesting is the η -pairing mechanism proposed by C.N.Yang in [2, 3, 4].

The one-dimensional Hubbard model has been known to be exactly solvable since the work of E.Lieb and F.Y.Wu of 1968 ^[5]. In their paper, a large set of eigenfunctions of the hamiltonian were found by using the nested Bethe Ansatz ^[6]. However, the issue of whether this set of eigenfunctions is actually *complete* has not been considered until recently. In a recent paper ^[7], we used the $SO(4)$ symmetry of the Hubbard model (which has been explored in [2, 3, 4, 8, 9]) to show that the Bethe Ansatz is not complete (see below for some comments on this result).

In this paper we will show that the $SO(4)$ structure can be used to extend the set of Bethe Ansatz eigenstates to a complete set of eigenstates of the one-dimensional Hubbard model. This result was first announced in [10]. Here we provide a detailed account of the derivations involved.

The Hubbard model describes electrons, which can hop along a one-dimensional lattice and which interact with coupling constant U if two of them occupy the same site. The hamiltonian of the Hubbard model on a periodic one-dimensional lattice of even, finite length L is given by (the small modifications in the potential term as compared to [5] are such that the $SO(4)$ symmetry becomes explicit, see [8, 2, 9, 3])

$$H = - \sum_{i=1}^L \sum_{\sigma=1,-1} (c_{i,\sigma}^\dagger c_{i+1,\sigma} + c_{i+1,\sigma}^\dagger c_{i,\sigma}) + U \sum_{i=1}^L (n_{i,1} - \frac{1}{2})(n_{i,-1} - \frac{1}{2}). \quad (1.1)$$

Here the $c_{i,\sigma}$ (σ takes the values ± 1) are canonical Fermi operators on the lattice ($i = 1, \dots, L$ labels the lattice sites), with anti-commutation relations given by $\{c_{i,\sigma}^\dagger, c_{j,\tau}\} = \delta_{i,j}\delta_{\sigma,\tau}$. They act in a Fock space with the pseudo vacuum $|0\rangle$ defined by $c_{i,\sigma}|0\rangle = 0$. The operator $n_{i,\sigma} = c_{i,\sigma}^\dagger c_{i,\sigma}$ is the number operator for electrons with spin σ on site i . U is the coupling constant and can be either positive (repulsive

case) or negative (attractive case). For later convenience we define

$$u = \frac{U}{2i}. \quad (1.2)$$

The analysis by E.Lieb and F.Y.Wu in [5] resulted in a large number of eigenstates of the hamiltonian, which are characterized by momenta k_i and rapidities Λ_α , where $i = 1, 2, \dots, N_e$ and $\alpha = 1, 2, \dots, M$ for an eigenstate with a total number of N spin-up and M spin-down electrons. Our convention throughout the paper will be

$$\begin{aligned} N &= \text{number of spin-up electrons} \\ M &= \text{number of spin-down electrons} \\ N_e &= N + M = \text{total number of electrons} . \end{aligned}$$

We will now discuss the Bethe wave functions wave functions in the form as given by Woynarovich [11], which is equivalent to the form found by Lieb and Wu. The nested Bethe Ansatz provides us with the following set of eigenstates with M spins down and N spins up :

$$\begin{aligned} |\Psi_{M,N}\rangle &= \sum_{1 \leq x_k \leq L} \psi_{-1, \dots, -1, 1, \dots, 1}(x_1, \dots, x_{N_e}) \prod_{j=1}^M c_{x_j, -1}^\dagger \prod_{i=M+1}^{N_e} c_{x_i, 1}^\dagger |0\rangle \\ &= \sum_{1 \leq x_k \leq L} \psi_{\sigma_1, \sigma_2, \dots, \sigma_{N_e}}(x_1, \dots, x_{N_e}) \prod_{j=1}^{N_e} c_{x_j, \sigma_j}^\dagger |0\rangle, \end{aligned} \quad (1.3)$$

where we have put $\sigma_1 = \dots = \sigma_M = -1$, $\sigma_{M+1} = \dots = \sigma_{N_e} = 1$.

The Bethe Ansatz wave functions explicitly depend on the relative ordering of the x_i . We represent this dependence by a permutation Q of N_e elements, which is such that $1 \leq x_{Q_1} \leq x_{Q_2} \leq \dots \leq x_{Q_{N_e}} \leq L$. In the sector Q the general Bethe wavefunction for M spins down and N spins up reads

$$\psi_{\sigma_1, \sigma_2, \dots, \sigma_{N_e}}(x_1, \dots, x_{N_e}) = \sum_{P \in S_{N_e}} \text{sgn}(Q) \text{sgn}(P) e^{i \sum_{j=1}^{N_e} k_{P_j} x_{Q_j}} \varphi(y_1, \dots, y_M | P). \quad (1.4)$$

The P -summation extends over all permutations of N_e elements and $\text{sgn}(\Pi)$ is the sign of the permutation Π ($\Pi = Q, P$). The amplitudes $\varphi(y_1, \dots, y_M | P)$ are of the form

$$\varphi(y_1, \dots, y_M | P) = \sum_{\pi \in S_M} A_\pi \prod_{l=1}^M F_P(\Lambda_{\pi_l}, y_l) \quad (1.5)$$

with

$$\begin{aligned}
F_P(\Lambda_j, y) &= \left(\prod_{i=1}^{y-1} \frac{e_-^{(j)}(P_i)}{e_+^{(j)}(P_i)} \right) \frac{1}{e_+^{(j)}(P_y)} \\
&= \left(\prod_{i=1}^{y-1} \frac{\sin(k_{P_i}) - \Lambda_j - \frac{U}{4i}}{\sin(k_{P_i}) - \Lambda_j + \frac{U}{4i}} \right) \left(\frac{1}{\sin(k_{P_y}) - \Lambda_j + \frac{U}{4i}} \right), \quad (1.6)
\end{aligned}$$

where we defined

$$e_{\pm}^{(j)}(i) = \sin(k_i) - \Lambda_j \pm \frac{u}{2} \quad (1.7)$$

and

$$\frac{A_{\pi}}{A_{(t,t+1)\pi}} = \frac{\Lambda_{\pi_{t+1}} - \Lambda_{\pi_t} - u}{\Lambda_{\pi_{t+1}} - \Lambda_{\pi_t} + u}. \quad (1.8)$$

By $\pi = (\pi_1, \pi_2, \dots, \pi_t, \pi_{t+1}, \dots, \pi_M)$ we denote a permutation of M elements (spin-down electrons) and $(t, t+1)\pi = (\pi_1, \pi_2, \dots, \pi_{t+1}, \pi_t, \dots, \pi_M)$. A solution of (1.8) is given by

$$A_{\pi} = \prod_{1 \leq l < k \leq M} \left(\frac{\Lambda_{\pi_l} - \Lambda_{\pi_k} + u}{\Lambda_{\pi_l} - \Lambda_{\pi_k}} \right). \quad (1.9)$$

The amplitudes $\varphi(y_1, \dots, y_M | P)$ depend on $\sigma_1, \dots, \sigma_{N_e}$ and on Q through the numbers $y_1, \dots, y_M$, which are defined to be the positions of the down spins among the spins in the series $\sigma_{Q_1}, \sigma_{Q_2}, \dots, \sigma_{Q_{N_e}}$ in increasing order, *i.e.*,

$$1 \leq y_1 < y_2 < y_3 < \dots < y_M \leq N_e. \quad (1.10)$$

For example, for one spin down and one spin up (and $\sigma_1 = -1, \sigma_2 = 1$) we have the two cases $y = 1$ (if the spin down is to the left, which holds in the $Q = (id)$ sector) and $y = 2$ (if the spin down is to the right, which holds for $Q = (21)$).

As we already indicated in (1.3), we will choose the notation such that the M down spins are at the positions $x_1, \dots, x_M$, *i.e.*, $\sigma_1 = \dots = \sigma_M = -1$ and $\sigma_{M+1} = \dots = \sigma_{N_e} = 1$.

We see that all solutions are characterised by N_e momenta $\{k_j | j=1, \dots, N_e\}$ of charged spinless excitations (holons), and M rapidities $\{\Lambda_k | k=1, \dots, M\}$ of spin waves (spinons).

Imposing periodic boundary conditions on the Bethe Ansatz wavefunctions leads to the following equations for the parameters k_i and Λ_α

$$e^{ik_j L} = \prod_{\alpha=1}^M \frac{\sin(k_j) - \Lambda_\alpha - \frac{U}{4i}}{\sin(k_j) - \Lambda_\alpha + \frac{U}{4i}}, \quad j = 1, 2, \dots, N_e,$$

$$\prod_{j=1}^{N_e} \frac{\sin(k_j) - \Lambda_\alpha - \frac{U}{4i}}{\sin(k_j) - \Lambda_\alpha + \frac{U}{4i}} = - \prod_{\beta=1}^M \frac{\Lambda_\beta - \Lambda_\alpha - \frac{U}{2i}}{\Lambda_\beta - \Lambda_\alpha + \frac{U}{2i}}, \quad \beta = 1, 2, \dots, M. \quad (1.11)$$

Energy and momentum, *i.e.*, the eigenvalues of the hamiltonian (1.1) and the logarithm of the translation operator, of the system in a state corresponding to a solution of (1.11) are

$$E_{N_e} = -2 \sum_{i=1}^{N_e} \cos(k_i) + \frac{U}{2} \left[\frac{L}{2} - N_e \right],$$

$$P = \sum_{i=1}^{N_e} k_i. \quad (1.12)$$

The second term in the expression for the energy is due to the shift of $n_{j,\sigma}$ by $\frac{1}{2}$ in (1.1).

Because of the antisymmetry of the product over $c^\dagger$'s under interchange of any two of them, the wave functions $\psi_{\sigma_1 \dots \sigma_{N_e}}(x_1 \dots x_{N_e})$ can be (and have been) chosen to be completely antisymmetric under the simultaneous exchange $x_k \longleftrightarrow x_j$ and $\sigma_k \longleftrightarrow \sigma_j$, *i.e.*,

$$\psi_{\sigma_1 \dots \sigma_j \dots \sigma_k \dots \sigma_{N_e}}(x_1 \dots x_j \dots x_k \dots x_{N_e}) = -\psi_{\sigma_1 \dots \sigma_k \dots \sigma_j \dots \sigma_{N_e}}(x_1 \dots x_k \dots x_j \dots x_{N_e}). \quad (1.13)$$

We now define 'regular' Bethe Ansatz states (for finite L), to be denoted by $|\psi_{M,N}\rangle$, by the properties that $N - M \geq 0$ (non-negative third component of the spin), $N_e \leq L$ (less than or equal to half filling), and that all Λ_α and all k_j are finite. Bethe Ansatz states with $N - M < 0$ and/or $N_e > L$ can be obtained from the regular Bethe Ansatz states by using simple symmetry operations, *i.e.*, reflection of the third component of the spin and particle/hole correspondence, which commute with the hamiltonian ^[12]. The model is invariant under spin-rotations, with the

corresponding $SU(2)$ generators given by

$$\zeta = \sum_{i=1}^L c_{i,1}^\dagger c_{i,-1}, \quad \zeta^\dagger = (\zeta)^\dagger, \quad \zeta_z = \frac{1}{2} \sum_{i=1}^L (n_{i,-1} - n_{i,1}). \quad (1.14)$$

(Note that ζ_z equals *minus* the third component of the total spin.) For even L the model has a second $SU(2)$ invariance, generated by ^[3]

$$\eta = \sum_{i=1}^L (-1)^i c_{i,1} c_{i,-1}, \quad \eta^\dagger = \sum_{i=1}^L (-1)^i c_{i,-1}^\dagger c_{i,1}^\dagger, \quad \eta_z = \frac{1}{2} \sum_{i=1}^L (n_{i,-1} + n_{i,1}) - \frac{L}{2}. \quad (1.15)$$

The raising operator $\eta^\dagger$ of this second $SU(2)$ creates a pair of two opposite-spin electrons on the same site, with momentum π . Combining the two $SU(2)$'s, which commute with the hamiltonian and with one another, leads to an $SO(4)$ invariance of the one-dimensional Hubbard model for even lattice lengths^[9, 3]. For a discussion of the theoretical and possible experimental consequences of the existence of this symmetry, which also exists in the Hubbard model in two or three dimensions, we refer to the papers [3, 4, 9, 13].

In a previous paper ^[7], we established the following remarkable property of the regular Bethe Ansatz eigenstates of the hamiltonian: they are all *lowest weight states* of the $SO(4)$ algebra (1.14), (1.15), *i.e.*,

$$\eta | \psi_{M,N} \rangle = 0, \quad \zeta | \psi_{M,N} \rangle = 0. \quad (1.16)$$

This shows that acting with the raising operators $\eta^\dagger$ and $\zeta^\dagger$ on $| \psi_{M,N} \rangle$ leads to new eigenstates of the hamiltonian, which are not in the regular Bethe Ansatz. In this way, every regular Bethe Ansatz state $| \psi_{M,N} \rangle$ is the lowest weight state in a multiplet of states, which form a representation of $SO(4)$. Since

$$\eta_z | \psi_{M,N} \rangle = \frac{1}{2} (N_e - L) | \psi_{M,N} \rangle, \quad \zeta_z | \psi_{M,N} \rangle = \frac{1}{2} (M - N) | \psi_{M,N} \rangle, \quad (1.17)$$

a state $| \psi_{M,N} \rangle$ has spin $\eta = \frac{1}{2} (L - N_e)$ with respect to the η -pairing $SU(2)$ algebra and spin $\zeta = \frac{1}{2} (N - M)$ with respect to the ζ $SU(2)$ algebra. The dimension of the corresponding $SO(4)$ multiplet is therefore given by

$$\dim_{M,N} = (2\eta + 1)(2\zeta + 1) = (L - N_e + 1)(N - M + 1). \quad (1.18)$$

The states in this multiplet are of the form

$$|\psi_{M,N}^{\alpha,\beta}\rangle = (\eta^\dagger)^\alpha (\zeta^\dagger)^\beta |\psi_{M,N}\rangle. \quad (1.19)$$

By symmetry, all the states that are highest or lowest weight states with respect to one of the $SU(2)$ algebras are again given by the Bethe Ansatz (although in general they are outside the *regular* Bethe Ansatz). All other states are *not* given by the Bethe Ansatz, which shows that for this model the Bethe Ansatz is not complete. The simplest example of a state that is outside the Bethe Ansatz is $\eta^\dagger |0\rangle$, which describes a single η -pair of momentum π . The fact that this state is outside the Bethe Ansatz was explicitly confirmed in [7].

It is the main purpose of this paper to show that, if one counts the number of eigenstates that are related to the regular Bethe Ansatz states by the $SO(4)$ symmetry, one finds 4^L , which is precisely the correct dimension of the Hilbert space of the model *. Thus we will conclude that the Bethe Ansatz together with the $SO(4)$ structure leads to a complete set of eigenstates of the one-dimensional Hubbard model.

The paper is organized as follows. In section 2 we discuss in some detail the nature of the solutions of the Bethe equations for the Hubbard model. In Appendix A we show that the so-called Λ and $k - \Lambda$ strings give wave functions that describe bound states. In section 3 we will then count eigenstates and prove completeness. An explicit construction of the $\frac{1}{2}(L+2)(L-1)$ solutions of the Bethe equations in the sector with one spin-up and one spin-down electron (as opposed to the somewhat indirect construction used in the general proof) is presented in Appendix B.

2 Solutions of the Bethe equations for the Hubbard model

Let us focus on the Bethe Equations (1.11), which express the fact that the Bethe Ansatz wavefunctions (1.4) satisfy periodic boundary conditions.

*There are 4 possible electron configurations per lattice site (spin up, spin down, both spin up and spin down, and empty site), thus the corresponding direct product Hilbert space is 4^L dimensional.

Counting regular Bethe Ansatz states means counting inequivalent solutions of the equations (1.11) while taking into account the ‘regularity conditions’ $N - M \geq 0$ and $N_e \leq L$. Following M.Takahashi ^[14], we will first distinguish different types of solutions $\{k_i, \Lambda_\alpha\}$ of (1.11). The idea is that for a solution $\{k_i, \Lambda_\alpha\}$, the set of all the k_i ’s and Λ_α ’s can be split into (three) different kinds of subsets (‘strings’), which are

1. a single real momentum k_i
2. m Λ_α ’s combine into a string-type configuration (‘ Λ -strings’); this includes the case $m = 1$, which is just a single real Λ_α [†]
3. $2m$ k_i ’s and m Λ_α ’s combine into a different string-type configuration (‘ k - Λ -strings’) [‡].

For large lattices ($1 \ll L$), almost all the string configurations are close to ‘idealized’ string-solutions where both the k ’s and the Λ ’s are assigned imaginary parts according to a ‘equal-spacing’ prescription ^[14]. For a Λ -string of length m the rapidities involved are

$$\Lambda_\alpha^{m,j} = \Lambda_\alpha^m - \frac{1}{2}(m+1-2j)u \quad \Lambda_\alpha^m \text{ real} \quad j = 1, 2, \dots, m. \quad (2.1)$$

The k ’s and Λ ’s involved in a k - Λ -string are

$$\begin{aligned} k_\alpha^1 &= \pi - \sin^{-1}(\Lambda_\alpha'^m - \frac{1}{2}mu) \\ k_\alpha^2 &= \sin^{-1}(\Lambda_\alpha'^m - \frac{1}{2}(m-2)u) \\ k_\alpha^3 &= \pi - k_\alpha^2 \\ k_\alpha^4 &= \sin^{-1}(\Lambda_\alpha'^m - \frac{1}{2}(m-4)u) \\ k_\alpha^5 &= \pi - k_\alpha^4 \\ &\dots \end{aligned}$$

[†]these correspond to bound states of spin waves (magnons)

[‡]the case $m = 1$ describes a ‘Cooper pair’ of electrons

$$\begin{aligned}
k_\alpha^{2m-2} &= \sin^{-1}(\Lambda_\alpha'^m + \frac{1}{2}(m-2)u) \\
k_\alpha^{2m-1} &= \pi - k_\alpha^{2m-2} \\
k_\alpha^{2m} &= \pi - \sin^{-1}(\Lambda_\alpha'^m + \frac{1}{2}mu) .
\end{aligned} \tag{2.2}$$

and

$$\Lambda_\alpha'^{m,j} = \Lambda_\alpha'^m - \frac{1}{2}(m+1-2j)u , \quad \Lambda_\alpha'^m \text{ real} \quad j = 1, 2, \dots, m . \tag{2.3}$$

(2.1) - (2.3) are valid up to exponential corrections of order $\mathcal{O}(\exp -\delta L)$, where δ is real and positive (and depends on the specific string under consideration). In Appendix A we discuss the wave-functions corresponding to some of the the string configurations 2. and 3. and show that they correspond to bound states.

Let us now consider a solution that splits into M_m Λ -strings of length m , M'_n k - Λ -strings of length n (containing $2n$ k_i 's and n Λ_α 's) and M_e additional single k_i 's. Clearly, we have

$$N_e = M_e + 2 \sum_{m=1}^{\infty} m M'_m , \quad M = \sum_{m=1}^{\infty} m(M_m + M'_m) . \tag{2.4}$$

How many solutions of this type exist?

The idea is that each of the strings in a solution can be characterised by the position of its center (a real number), which we denote as in (2.1) by Λ_α^m , $\alpha = 1, 2, \dots, M_m$, for the length- m Λ -strings, by $\Lambda_\alpha'^m$, $\alpha = 1, 2, \dots, M'_m$ for the length- m k - Λ -strings (as in (2.3)) and which is simply equal to k_j for the unpaired momenta k_j , $j = 1, 2, \dots, M_e$. Because of the periodic boundary conditions, these parameters will have to be chosen from a discrete set.

Following [14], we now write the following equations for the parameters k_j , Λ_α^m and $\Lambda_\alpha'^m$. They follow from (1.11) and the form of the 'idealized' string-solutions which we discussed above (we write $M' = \sum_{m=1}^{\infty} m M'_m$)

$$\begin{aligned}
k_j L &= 2\pi I_j - \sum_{n=1}^{\infty} \sum_{\alpha=1}^{M_n} \theta \left(\frac{\sin k_j - \Lambda_\alpha^n}{nU} \right) - \sum_{n=1}^{\infty} \sum_{\alpha=1}^{M'_n} \theta \left(\frac{\sin k_j - \Lambda_\alpha'^n}{nU} \right) , \\
\sum_{j=1}^{N_e-2M'} \theta \left(\frac{\Lambda_\alpha^n - \sin k_j}{nU} \right) &= 2\pi J_\alpha^n + \sum_{m=1}^{\infty} \sum_{\beta=1}^{M_m} \Theta_{nm} \left(\frac{\Lambda_\alpha^n - \Lambda_\beta^m}{U} \right) ,
\end{aligned}$$

$$\begin{aligned}
& L \left[\sin^{-1} \left(\Lambda'_\alpha + in \frac{U}{4} \right) + \sin^{-1} \left(\Lambda'_\alpha - in \frac{U}{4} \right) \right] \\
&= 2\pi J'_\alpha + \sum_{j=1}^{N_e-2M'} \theta \left(\frac{\Lambda'_\alpha - \sin k_j}{nU} \right) + \sum_{m=1}^{\infty} \sum_{\beta=1}^{M'_m} \Theta_{nm} \left(\frac{\Lambda'_\alpha - \Lambda'_\beta}{U} \right), \quad (2.5)
\end{aligned}$$

where

$$\begin{aligned}
\theta(x) &= 2 \tan^{-1}(4x) \\
\Theta_{nm}(x) &= \begin{cases} \theta \left(\frac{4x}{|n-m|} \right) + 2\theta \left(\frac{4x}{|n-m|+2} \right) + \dots + 2\theta \left(\frac{4x}{n+m-2} \right) + \theta \left(\frac{4x}{n+m} \right) & \text{for } n \neq m \\ 2\theta \left(\frac{2x}{1} \right) + \dots + 2\theta \left(\frac{2x}{n-1} \right) + \theta \left(\frac{2x}{n} \right) & \text{for } n = m \end{cases} \quad (2.6)
\end{aligned}$$

The I_j , J_α^n and J'_α are integer or half-odd-integer according to the following prescriptions: I_j is integer (half-odd-integer) if $\sum_m (M_m + M'_m)$ is even (odd); the J_α^n are integer (half-odd-integer) if $(N_e - M_n)$ is odd (even); the J'_α are integer (half-odd-integer) if $(L - (N_e - M'_n))$ is odd (even). According to [14], we have the following inequalities

$$\begin{aligned}
|J'_\alpha| &\leq \frac{1}{2} (N_e - 2M' - \sum_{m=1}^{\infty} t_{nm} M_m - 1), \\
|J_\alpha^n| &\leq \frac{1}{2} (L - N_e + 2M' - \sum_{m=1}^{\infty} t_{nm} M'_m - 1), \\
0 < I_j &\leq L, \quad (2.7)
\end{aligned}$$

where $t_{nm} = 2 \text{Min}(n, m) - \delta_{nm}$.

We will now make the standard assumption that, in order to enumerate the different solutions of the system (2.5), it is sufficient to enumerate all possible sets of *non-repeating* (half-odd) integers I_j , J_α^n and J'_α , satisfying (2.7).

[This assumption mimics the similar assumption which is usually made for the spin- $\frac{1}{2}$ Heisenberg XXX model [15, 16]. It is known, however[§], that the actual distribution of the different types of solutions can be different from the one implied

[§]This fact was actually already noticed in the original paper by Bethe, [17].

by this counting. We carefully studied this phenomenon in a recent paper [18], where we give a detailed discussion of the two-magnon sector of the XXX model. We explicitly show that the deviations from the above assumption can be viewed as a ‘redistribution phenomenon’, which does not affect the total number of Bethe Ansatz states. For the Hubbard model we have a similar situation, which gets complicated further by the fact that there is a free coupling constant U in the model. In Appendix B we analyze in detail the $N = 1$, $M = 1$ sector of the Hubbard model. Although we do find U -dependent redistributions among different types of solutions, we find agreement with the predictions based on the ‘ideal’ string assumption for the total number of states in this sector, which is $\frac{1}{2}(L-1)(L+2)$ for the L -site model.]

From (2.7) we read off that the numbers of allowed values for the (half-odd-)integers corresponding to each of the fundamental strings are

1. L for a free k_i
2. $N_e - 2M' - \sum_{m=1}^{\infty} t_{nm} M_m$ for a Λ -string of length n
3. $L - N_e + 2M' - \sum_{m=1}^{\infty} t_{nm} M'_m$ for a k - Λ -string of length n .

The total number of ways to choose the (half-odd-)integers in a solution with multiplicities M_e , M_m and M'_m is therefore given by (remember that the integers are assumed to be non-repeating)

$$n(M_e, \{M_m\}, \{M'_m\}) = \binom{L}{M_e} \prod_{n=1}^{\infty} \binom{N_e - 2M' - \sum_{m=1}^{\infty} t_{nm} M_m}{M_n} \times \\ \times \prod_{n=1}^{\infty} \binom{L - N_e + 2M' - \sum_{m=1}^{\infty} t_{nm} M'_m}{M'_n}. \quad (2.8)$$

The total number of solutions of (1.11) with given numbers N and M is now obtained by summing $n(M_e, \{M_m\}, \{M'_m\})$ over all the M_e , M_m and M'_m , under the constraints (2.4).

Every solution to (1.11) gives us a regular Bethe Ansatz state, which comes with an entire multiplet of eigenstates of the hamiltonian, the dimension $\dim_{M,N}$ of which is given in (1.18). The full number of eigenstates that are obtained from the Bethe

Ansatz and the $SO(4)$ symmetry is therefore given by

$$\# \text{ (eigenstates)} = \sum_{\substack{M \geq 0 \\ N-M \geq 0 \\ N+M \leq L}} \sum_{\substack{N \geq 0 \\ N+M=M_e+2 \sum_{m=1}^{\infty} m M'_m \\ N-2M=M_e-2 \sum_{m=1}^{\infty} m M_m}} \left[\sum_{M_e=0}^{\infty} \sum_{M_m=0}^{\infty} \sum_{M'_m=0}^{\infty} n(M_e, \{M_m\}, \{M'_m\}) \right] \dim_{M,N}. \quad (2.9)$$

The counting of the eigenstates that are obtained from the $SO(4)$ extended nested Bethe Ansatz has thus been reduced to a purely algebraic problem, which we will solve in the next section.

3 Counting eigenstates

In this section we will prove that for general even L the sum in (2.9) equals 4^L . This will prove completeness. Before we come to that, we show the examples of the 2-site and 4-site models. The 2-site model ($L = 2$) was discussed in [7], where we presented the explicit form of a complete set of $4^2 = 16$ eigenstates of the hamiltonian. In Table 1 we show how the counting presented in section 3 works out in this case.

M_e	M_1	M'_1	M	N	n	$\dim_{M,N}$	$\#(\text{states})$
0	0	0	0	0	1	3	3
1	0	0	0	1	2	4	8
2	0	0	0	2	1	3	3
2	1	0	1	1	1	1	1
0	0	1	1	1	1	1	1
							16

Table 1: $L = 2$. n denotes the number of regular Bethe Ansatz states of a given type. There are a total number of $16 = 4^2$ eigenstates of the hamiltonian.

The total number of 16 states splits into 2 singlets, 2 triplets and 2 quadruplets of $SO(4)$. The ground state is the singlet with $M_1 = 1$ for the case $U > 0$ and the

M_e	M_1	M_2	M'_1	M'_2	M	N	n	$\dim_{M,N}$	#(states)
0	0	0	0	0	0	0	1	5	5
1	0	0	0	0	0	1	4	8	32
2	0	0	0	0	0	2	6	9	54
3	0	0	0	0	0	3	4	8	32
4	0	0	0	0	0	4	1	5	5
2	1	0	0	0	1	1	6	3	18
0	0	0	1	0	1	1	3	3	9
3	1	0	0	0	1	2	8	4	32
1	0	0	1	0	1	2	8	4	32
4	2	0	0	0	2	2	1	1	1
4	0	1	0	0	2	2	1	1	1
2	1	0	1	0	2	2	6	1	6
0	0	0	2	0	2	2	1	1	1
0	0	0	0	1	2	2	1	1	1
4	1	0	0	0	1	3	3	3	9
2	0	0	1	0	1	3	6	3	18
									256

Table 2: $L = 4$. There are 60 regular Bethe Ansatz states, which, when weighted with the correct $SO(4)$ multiplicities, give a total of $256 = 4^4$ eigenstates of the hamiltonian.

singlet with $M'_1 = 1$ for the case $U < 0$. In both cases it is a bound state of one spin up and one spin down electron with energy $E_0 = -\sqrt{U^2/4 + 16}$. The counting for the 4-site model ($L = 4$) is presented in Table 2, where we show how the total number of $4^4 = 256$ is obtained. (Notice that the total number of regular Bethe Ansatz states is only 60.)

We now turn to the proof that for general (even) L the sum in (2.9) equals 4^L . We will split this proof into two steps as follows. In the first step we will prove the following two identities

$$\sum_{\substack{M_1, M_2, \dots = 0 \\ \sum_{m=1}^{\infty} m M_m = M}}^{\infty} \prod_{n=1}^{\infty} \binom{N - \sum_m t_{nm} M_m}{M_n} = \binom{N}{M} - \binom{N}{M-1} \quad (3.1)$$

and

$$\sum_{M=0}^{[N/2]} \left(\binom{N}{M} - \binom{N}{M-1} \right) (N - 2M + 1) = 2^N. \quad (3.2)$$

For later convenience we define

$$P_n = N - \sum_{m=1}^{\infty} t_{nm} M_m, \quad (3.3)$$

and

$$n(\{M_m\}) = \prod_{n=1}^{\infty} \binom{P_n}{M_n}, \quad (3.4)$$

where $t_{nm} = 2 \text{Min}(n, m) - \delta_{nm}$ as before. In the second step we will then use identities (3.1) and (3.2) to perform the summation in (2.9).

The auxiliary identities (3.1) and (3.2) have a natural interpretation in the context of the spin-1/2 Heisenberg XXX model [16]. The equation (3.1) gives the total number of regular Bethe Ansatz states (defined by $M \leq [N/2]$) with M overturned spins in the XXX model on a lattice of length N . The second formula shows that the total number of states obtained by combining the regular Bethe Ansatz with

the $SU(2)$ structure equals 2^N , which is the dimension of the Hilbert space of the XXX model. These relations thus establish the completeness of the $SU(2)$ extended Bethe Ansatz for the XXX model.

The fact that identities that have their origin in the XXX model play a role here should not come as a surprise. Indeed, our method of solution of the Hubbard model is the *nested* Bethe Ansatz. The solutions to the Bethe Ansatz are specified by two sets $\{k_j\}$ and $\{\Lambda_\alpha\}$ of spectral parameters. The k_j 's are momenta of charge density waves, whereas the Λ_α 's, which describe the 'nesting' of the Bethe Ansatz, are rapidities of spin density waves of the type encountered in the Heisenberg XXX model. This should make clear that the second stage of the nested Bethe Ansatz for the Hubbard model is really a spin-problem, which is very similar to the Bethe Ansatz analysis of the Heisenberg XXX model. Our two-step procedure for performing the summation is natural from the point of view of the nesting: in the first step we sum over the spin degrees of freedom, and in the second step we then sum over the charge degrees of freedom as well.

STEP 1.

Let us now explain how the equation (3.1) can be derived. In the first step, one simply solves for $M_1 = M - \sum_{m=2}^{\infty} m M_m$ and substitutes this back into the left hand side of (3.1). Using this value for M_1 , the quantities P_n reduce to

$$\begin{aligned} P_1 &= N - \sum_{m=1}^{\infty} t_{1m} M_m = N - M + \sum_{m=3}^{\infty} (m-2) M_m, \\ P_n &= N - \sum_{m=1}^{\infty} t_{nm} M_m = N - 2M + M_n + 2 \sum_{m=n+1}^{\infty} (m-n) M_m. \end{aligned} \quad (3.5)$$

Let us now consider the summation over M_2 . Although our summand in the left hand side of (3.1) has the form of an infinite product, only two of the factors contain the variable M_2 . Singling these out, one finds that the summation over M_2 is as follows

$$\Omega_2 = \sum_{M_2=0}^{\infty} \left(\frac{N - 2M + M_2 + 2 \sum_{m=3}^{\infty} (m-2) M_m}{M_2} \right) \times$$

$$\times \binom{N - M + \sum_{m=3}^{\infty} (m-2)M_m}{M - \sum_{m=2}^{\infty} m M_m} . \quad (3.6)$$

In order to perform this summation we will make use of the identity

$$\sum_{\alpha=0}^{\infty} \binom{B + \alpha}{\alpha} x^{\alpha} = (1 - x)^{-1-B} , \quad (3.7)$$

which can easily be proved by induction. As a simple consequence, we have

$$(1 - x^2)^{-1-\omega} (1 + x)^{\eta} |_{x^A} = \sum_{\alpha=0}^{\infty} \binom{\omega + \alpha}{\alpha} \binom{\eta}{A - 2\alpha} , \quad (3.8)$$

where the notation $|_{x^A}$ in the left hand side means that we single out the coefficient of the power x^A . The right hand side of (3.6) is of the same form as (3.8) and we find

$$\begin{aligned} \Omega_2 &= (1 - x^2)^{-1-[N-2M+2(M_3+2M_4+\dots)]} (1 + x)^{N-M+(M_3+2M_4+\dots)} |_{x^{M-3M_3-4M_4-\dots}} \\ &= (1 + x)^{N-M} (1 - x^2)^{-N+2M-1} \prod_{n=3}^{\infty} (\mathcal{Z}_n^{(0)})^{M_n} |_{x^M} \\ &= \frac{1}{2\pi i} \oint \frac{dx}{x^{M+1}} (1 + x)^{N-M} (1 - x^2)^{-N+2M-1} \prod_{n=3}^{\infty} (\mathcal{Z}_n^{(0)})^{M_n} , \end{aligned} \quad (3.9)$$

where

$$\mathcal{Z}_n^{(0)} = \frac{x^n}{(1 - x)^{2(n-2)} (1 + x)^{n-2}} . \quad (3.10)$$

In the last line of (3.9) we extracted the coefficient at x^M by performing a contour integral around the origin $x = 0$. After performing the M_2 summation (3.1) now reads

$$\sum_{\substack{M_1, M_2, \dots = 0 \\ \sum_{m=1}^{\infty} m M_m = M}}^{\infty} n(\{M_m\}) = \frac{1}{2\pi i} \oint \frac{dx}{x^{M+1}} A(x) , \quad (3.11)$$

where

$$A(x) = (1+x)^{N-M}(1-x^2)^{-1-N+2M} \times \\ \times \sum_{M_3, M_4, \dots=0}^{\infty} \prod_{n=3}^{\infty} \binom{N-2M+M_n+2\sum_{m=n+1}^{\infty}(m-n)M_m}{M_n} \prod_{l=3}^{\infty} (\mathcal{Z}_l^{(0)})^{M_l}. \quad (3.12)$$

The summation over M_3 is given by

$$\Omega_3 = \sum_{M_3=0}^{\infty} \binom{N-2M+M_3+2\sum_{n=4}^{\infty}(n-3)M_n}{M_3} (\mathcal{Z}_3^{(0)})^{M_3} \\ = (1-\mathcal{Z}_3^{(0)})^{-1-N+2M-2\sum_{n=4}^{\infty}(n-3)M_n}. \quad (3.13)$$

At this point, the full expression for $A(x)$ has been reduced to

$$A(x) = (1+x)^{N-M}(1-x^2)^{-1-N+2M} (1-\mathcal{Z}_3^{(0)})^{-1-N+2M} \times \\ \times \sum_{M_4, M_5, \dots=0}^{\infty} \prod_{n=4}^{\infty} \binom{N-2M+M_n+2\sum_{m=n+1}^{\infty}(m-n)M_m}{M_n} \prod_{l=4}^{\infty} (\mathcal{Z}_l^{(1)})^{M_l}, \quad (3.14)$$

where $\mathcal{Z}_n^{(1)}$ are defined through

$$\mathcal{Z}_n^{(1)} = \frac{\mathcal{Z}_n^{(0)}}{(1-\mathcal{Z}_3^{(0)})^{2(n-3)}}. \quad (3.15)$$

From the above it is now clear, that the sum with respect to M_4 and all M_n with $n > 4$ has the same structure as the sum with respect to M_3 . Thus the final result, after performing all summations, will look like

$$A(x) = (1+x)^{N-M} F(x)^{-1-N+2M}, \quad (3.16)$$

where

$$F(x) = (1-x^2) \prod_{m=3}^{\infty} (1-\mathcal{Z}_m^{(m-3)}) \quad (3.17)$$

and we have the iteration formula

$$\mathcal{Z}_n^{(m)} = \frac{\mathcal{Z}_n^{(m-1)}}{\left(1 - \mathcal{Z}_{m+2}^{(m-1)}\right)^{2(n-m-2)}}. \quad (3.18)$$

Our task is now to find a closed expression for $F(x)$ by exploiting this relation. We define

$$U_2 = x^{-2}, \quad U_m = \frac{1}{\mathcal{Z}_m^{(m-3)}}, \quad m \geq 3, \quad (3.19)$$

so that $F(x)$ can be written as

$$F(x) = \prod_{m=2}^{\infty} \left(1 - \frac{1}{U_m}\right). \quad (3.20)$$

We now claim that the functions $U_m(x)$ satisfy the following recursion relation, to be denoted by $RR I_p$

$$RR I_p : \quad (U_{p+3} - 1)^2 = U_{p+4} U_{p+2}, \quad p \geq 0. \quad (3.21)$$

Together with the initial conditions

$$U_2 = x^{-2}, \quad U_3 = \frac{(1-x)^2(1+x)}{x^3} \quad (3.22)$$

these relations completely fix the functions $U_m(x)$ and thereby the function $F(x)$.

In order to prove the recursion relation (3.21), we first give a second recursion relation, which involves some of the other $\mathcal{Z}$'s and which we shall denote by $RR II_p$

$$RR II_p : \quad \frac{\mathcal{Z}_{n+1}^{(p)}}{\mathcal{Z}_n^{(p)}} = \frac{U_{p+2}}{U_{p+3}}, \quad p \geq 0, \quad n \geq p+3. \quad (3.23)$$

Let us now show that the validity of both recursion relations can be proved by induction. We start at the point where we have $\mathcal{Z}_n^{(0)}$, which is defined by (3.10), and U_2 and U_3 as above in (3.22). One easily checks that $RR II_{p=0}$ is valid. Using (3.18) for the definition of $U_4 = 1/\mathcal{Z}_4^{(1)}$ and $RR II_{p=0}$, one proves $RR I_{p=0}$. This

establishes the validity of both $RR I_{p=0}$ and $RR II_{p=0}$, which is the starting point for the proof by induction.

Let us now assume that we have proved the validity of both $RR I_p$ and $RR II_p$ for some given p . By using this induction assumption and the definition (3.18), one then proves the relation $RR II_{p+1}$ (3.23). After that, by using the definition (3.18) and $RR II_{p+1}$, one then proves $RR I_{p+1}$. This completes the induction step. We may thus conclude that the relations $RR I_p$ and $RR II_p$ are valid for all $p \geq 0$.

One easily checks that the expressions

$$U_j = \left(\frac{a(x)^{j+1} - a(x)^{-j-1}}{a(x) - a(x)^{-1}} \right)^2 \quad (3.24)$$

with

$$a(x) = \frac{1}{2} \left(\sqrt{\frac{1-3x}{x}} + \sqrt{\frac{1+x}{x}} \right) \quad (3.25)$$

satisfy the recursion relations (3.21) and the initial conditions (3.22). The function $F^2(x)$ is now expressed as a convergent product

$$F^2(x) = \prod_{m=2}^{\infty} \frac{(U_m - 1)^2}{U_m^2} = \prod_{m=2}^{\infty} \frac{U_{m+1} U_{m-1}}{U_m^2} = \lim_{l \rightarrow \infty} \frac{U_1 U_{l+1}}{U_2 U_l} = a(x)^2 x(x+1), \quad (3.26)$$

where we defined $U_1(x) = \frac{x+1}{x}$, in accord with (3.21), and where we used (3.21) in the second equality. This brings us to the following representation of the number of regular Bethe Ansatz states with M overturned spins (using (3.1), (3.11), (3.12), (3.16), (3.20), (3.25) and (3.26))

$$\begin{aligned} \sum_{\substack{M_m=0 \\ M=\sum_{m=1}^{\infty} m M_m}}^{\infty} n(\{M_m\}) &= \\ &= \frac{1}{2\pi i} \oint \frac{dx}{x^{M+1}} (1+x)^{N-M} \left(\frac{(1+x) + \sqrt{(1+x)(1-3x)}}{2} \right)^{2M-N-1} \\ &= \frac{1}{2\pi i} \oint \frac{2dy}{y^{M+1}} (2(1+y))^{N-M} \left(1+y + \sqrt{1-y^2} \right)^{2M-N-1}, \end{aligned} \quad (3.27)$$

where the contour is a small circle around the origin and we used the substitution $y = \frac{2x}{1-x}$. Calling $y^{-1} = \cosh \phi$, the integral reduces to $I_+ - I_-$, where

$$I_{\pm} = \frac{1}{2\pi} \int_0^{2\pi} d\varphi e^{-\phi(N-M \mp 1)} (1 + e^{\phi})^{N-1} = \binom{N-1}{N-M \mp 1} \quad (3.28)$$

where we wrote $\phi = \Lambda - i\varphi$, with $\Lambda \rightarrow \infty$. This finally establishes the result (3.1) since

$$\begin{aligned} \sum_{\substack{M_m=0 \\ M=\sum_{m=1}^{\infty} m M_m}}^{\infty} n(\{M_m\}) &= \binom{N-1}{N-M-1} - \binom{N-1}{N-M+1} \\ &= \binom{N}{M} - \binom{N}{M-1}. \end{aligned} \quad (3.29)$$

We still have to prove (3.2), which can be done as follows :

$$\begin{aligned} \sum_{M=0}^{[N/2]} \left(\binom{N}{M} - \binom{N}{M-1} \right) (N - 2M + 1) &= \\ &= \sum_{M=0}^{[N/2]} \binom{N}{M} (N - 2M + 1) - \sum_{M=0}^{[N/2]-1} \binom{N}{M} (N - 2M - 1) \\ &= 2 \sum_{M=0}^{[N/2]-1} \binom{N}{M} + \binom{N}{\lfloor \frac{N}{2} \rfloor} \left(1 + N - 2 \left\lfloor \frac{N}{2} \right\rfloor \right) \\ &= \sum_{M=0}^{[N/2]-1} \binom{N}{M} + \binom{N}{\lfloor \frac{N}{2} \rfloor} \left(1 + N - 2 \left\lfloor \frac{N}{2} \right\rfloor \right) + \sum_{N-[N/2-1]}^N \binom{N}{M} \\ &= \sum_{M=0}^N \binom{N}{M} = 2^N. \end{aligned} \quad (3.30)$$

This completes the proof of equation (3.2).

STEP 2.

The total number of states that are obtained from the SO(4) extended Bethe Ansatz for the Hubbard model is given by (2.8) and (2.9). The summations over the multiplicities M_m and over the difference $N - M$ in the summation (2.9) are precisely of the type (3.1) and (3.2), respectively, if we substitute $M \rightarrow \frac{1}{2}(M_e - N + M)$ and $N \rightarrow M_e$. (Under these summations the total number of electrons N_e is kept fixed.) The summation that remains after this ‘spin summation’ is

$$\begin{aligned} \# \text{ (eigenstates)} &= \sum_{N_e=0}^L (L - N_e + 1) \\ &\times \left[\sum_{M_e=0}^{N_e} \sum_{M'_m=0}^{\infty} 2^{M_e} \binom{L}{M_e} \prod_{n=1}^{\infty} \binom{L - N_e + \sum_{m=1}^{\infty} (2m - t_{nm}) M'_m}{M'_n} \right], \end{aligned} \quad (3.31)$$

where as before $t_{nm} = 2 \text{Min}(n, m) - \delta_{nm}$. In the next step we perform the summation with respect to all M'_n ’s, using a similar kind of ‘summation device’ as in STEP 1. As a consequence of (3.7) we have

$$(1 - x^2)^{-1-B} (1 + 2x)^L = \sum_{M'_1=0}^{\infty} \sum_{p=0}^L \binom{B + M'_1}{M'_1} \binom{L}{p} 2^p x^{2M'_1+p}, \quad (3.32)$$

and therefore

$$\frac{1}{2\pi i} \oint \frac{dx}{x^{\gamma+1}} (1 - x^2)^{-1-B} (1 + 2x)^L = \sum_{M'_1=0}^{\infty} \binom{B + M'_1}{M'_1} \binom{L}{\gamma - 2M'_1} 2^{\gamma-2M'_1}. \quad (3.33)$$

The integration is along a small contour around zero. Defining $E = L - N_e$, $\gamma = N_e - 2 \sum_{m=2}^{\infty} m M'_m$ and $B = E + 2 \sum_{m=2}^{\infty} (m - 1) M'_m$, the r.h.s. of (3.33) becomes the summation over M'_1 in (3.31), if we solve the constraint in the sum in (3.31) for $M_e = N_e - 2 \sum_{m=1}^{\infty} m M'_m$. Using (3.33) in (3.31) we then obtain the following expression for the number of eigenstates :

$$\# \text{ (eigenstates)} = \frac{1}{2\pi i} \oint \frac{dx}{x^{L+1}(1 - x^2)} (1 + 2x)^L \sum_{E=0}^L (E + 1) \frac{x^E}{(1 - x^2)^E} F(x), \quad (3.34)$$

where

$$F(x) = \sum_{\substack{M'_m=0 \\ m \geq 2}}^{\infty} \prod_{n=2}^{\infty} \binom{E + M'_n + 2 \sum_{m=n+1}^{\infty} (m-n) M'_m}{M'_n} \prod_{m=2}^{\infty} (\mathcal{Z}_m^{(0)})^{M'_m}, \quad (3.35)$$

and

$$\mathcal{Z}_m^{(0)} = \frac{x^{2m}}{(1-x^2)^{2(m-1)}}. \quad (3.36)$$

The summations over $M'_2, M'_3, \dots$ have precisely the form of the l.h.s. of (3.7) and can thus be performed easily. The result is

$$F(x) = \prod_{m=2}^{\infty} (1 - \mathcal{Z}_m^{(m-2)})^{-1-E}, \quad (3.37)$$

where

$$\mathcal{Z}_m^{(p)} = \frac{\mathcal{Z}_m^{(p-1)}}{(1 - \mathcal{Z}_{p+1}^{(p-1)})^{2(m-p-1)}}. \quad (3.38)$$

It can be shown along the lines given in STEP 1, that the quantities $U_m = \frac{1}{\mathcal{Z}_m^{(m-2)}}$ obey the recursion relation

$$(U_{p+2} - 1)^2 = U_{p+3} U_{p+1}, \quad p \geq 0 \quad (3.39)$$

with initial conditions

$$U_1 = x^{-2}, \quad U_2 = \frac{(1-x^2)^2}{x^4}. \quad (3.40)$$

Equation (3.23) is replaced by

$$\frac{\mathcal{Z}_{n+1}^{(p)}}{\mathcal{Z}_n^{(p)}} = \frac{U_{p+1}}{U_{p+2}}, \quad p \geq 0, \quad n \geq p+2. \quad (3.41)$$

Equation (3.34) now can be written as

$$\# \text{ (eigenstates)} = \frac{1}{2\pi i} \oint \frac{dx}{x^{L+1}} (1+2x)^L \sum_{E=0}^L (E+1) x^E [f(x)]^{-E-1}, \quad (3.42)$$

where

$$f(x) = \prod_{l=1}^{\infty} (1 - U_l^{-1}). \quad (3.43)$$

The solution of the recursion relation (3.39) is again of the form (3.24), i.e., $U_j = \left(\frac{a(x)^{j+1} - a(x)^{-j-1}}{a(x) - a(x)^{-1}} \right)^2$, where now

$$a(x) = \frac{1}{2x} + \sqrt{\frac{1}{4x^2} - 1} \quad (3.44)$$

due to the new initial conditions (3.40). Insertion of the resulting expression for U_l into (3.43) leads to the following result for the function $f(x)$:

$$2f(x) = 1 + \sqrt{1 - 4x^2} = 2x a(x). \quad (3.45)$$

Equation (3.42) can now be rewritten as

$$\# \text{ (eigenstates)} = \sum_{E=0}^L (E+1) I(E), \quad (3.46)$$

where

$$I(E) = \frac{1}{2\pi i} \oint d\left(-\frac{1}{x}\right) \left(\frac{1}{x} + 2\right)^L [a(x)]^{-E-1}. \quad (3.47)$$

The contour integration can be worked out as in section 2. Defining $\alpha = \Lambda - i\varphi$ with $\Lambda \gg 1$ and substituting $x = \frac{1}{e^\alpha + e^{-\alpha}}$ we obtain

$$I(E) = I_+(E) - I_-(E), \quad (3.48)$$

where

$$I_{\pm}(E) = \frac{1}{2\pi} \int_0^{2\pi} d\varphi e^{\pm\alpha} e^{-(1+E)\alpha} \left(e^{\frac{\alpha}{2}} + e^{-\frac{\alpha}{2}} \right)^{2L}. \quad (3.49)$$

Expanding

$$\left(e^{\frac{\alpha}{2}} + e^{-\frac{\alpha}{2}}\right)^{2L} = \sum_{p=0}^{2L} \binom{2L}{p} e^{\alpha(L-p)} \quad (3.50)$$

and then using

$$\frac{1}{2\pi} \int_0^{2\pi} d\varphi e^{\pm i(n\varphi)} = \delta_{n,0} \quad (3.51)$$

in the resulting expression, we find that

$$I_+(E) = \binom{2L}{L-E}, \quad I_-(E) = \binom{2L}{L-E-2}. \quad (3.52)$$

Plugging these results into (3.48) and then (3.46) we are left with only a single summation

$$\#(\text{eigenstates}) = \sum_{E=0}^L (E+1) \left\{ \binom{2L}{L-E} - \binom{2L}{L-E-2} \right\}. \quad (3.53)$$

This summation can be performed the same way as (3.30) and we finally obtain the desired result

$$\#(\text{eigenstates}) = 4^L. \quad (3.54)$$

This concludes our two-step evaluation of the sum (2.9).

Using the above, we can obtain a closed expression for the number of regular Bethe Ansatz states for given numbers M and N of spin-down and spin-up electrons:

$$\begin{aligned} \sum_{M_e=0}^{\infty} \sum_{M_m=0}^{\infty} \sum_{M'_m=0}^{\infty} n(M_e, \{M_m\}, \{M'_m\}) = \\ N_e = M_e + 2 \sum m M'_m \\ M = \sum m(M_m + M'_m) \end{aligned} \quad \left(\binom{L}{N} \left(\binom{L}{M} + \binom{L}{M-2} \right) - \left(\binom{L}{N+1} + \binom{L}{N-1} \right) \binom{L}{M-1} \right). \quad (3.55)$$

This formula is the close analogue of the result (3.1) for the XXX Heisenberg model.

We repeat once more our conclusion, which is that the combination of the nested Bethe Ansatz with the $SO(4)$ symmetry of the one-dimensional Hubbard model leads to a complete set of 4^L independent eigenstates.

It is a pleasure to thank C.N. Yang for proposing the ideas worked out in this paper. We thank S. Dasmahapatra for stimulating discussions. This work was supported by NSF Grant PHY- 9107261.

References

- [1] see for example the *Dialogue on the Theory of High T_c* by P.W. Anderson and R. Schrieffer in Physics Today (June 1991), 54-61
- [2] C.N. Yang, Phys.Rev.Lett. **63** (1989) 2144
- [3] C.N. Yang and S. Zhang, Mod.Phys.Lett. **B4** (1990) 759
- [4] C.N. Yang, preprint ITP-SB-91-38
- [5] E. Lieb and F.Y. Wu, Phys.Rev.Lett. **20** (1968) 1445
- [6] C.N. Yang, Phys.Rev.Lett. **19** (1967) 1315
- [7] F.H.L. Eßler, V.E. Korepin and K. Schoutens, *New Eigenstates of the 1-dimensional Hubbard Model*, preprint ITP-SB-91-15, June 1991 Nucl.Phys. **B**, to be published
- [8] O.J. Heilmann and E.H. Lieb, Annals of the New York Acad. of Sci. **172** (1971) 583; E.H. Lieb, Phys.Rev.Lett. **62** (1989) 1201
- [9] I. Affleck in talk given at the Nato Advanced Study Institute on *Physics, Geometry and Topology*, Banff, August 1989; M. Pernici, Europhys.Lett. **12** (1990) 75;
- [10] F.H.L. Eßler, V.E. Korepin and K. Schoutens, *Complete Solution of the One-dimensional Hubbard Model*, preprint ITP-SB-91-30 (July 1991)
- [11] F. Woynarovich, Jour. Phys. **C15** (1982) 85

- [12] B. Sutherland, *Lecture Notes in Physics*, vol. **242**, (Springer Verlag, Berlin, 1985) p. 1
- [13] S. Zhang, Phys.Rev.Lett. **65** (1990) 120
- [14] M. Takahashi, Prog.Theor.Phys. **47** (1972) 69
- [15] L.D. Faddeev and L. Takhtajan, Phys.Lett. **85A** (1981) 375; *ibid.*, Zap.Nauch.-Semin. LOMI, vol. **109** (1981) 134; *ibid.*, *Lecture Notes in Physics*, vol. **242**, (Springer Verlag, Berlin, 1985) p. 158, 175
- [16] M. Takahashi, Prog.Theor.Phys. **46** (1971) 401
- [17] H. Bethe, Z.Phys. **71** (1931) 205
- [18] F.H.L. Eßler, V.E. Korepin and K. Schoutens, *Fine Structure of the Bethe Ansatz for the spin- $\frac{1}{2}$ Heisenberg XXX Model* preprint ITP-SB-91-48 (Nov. 1991)

A Bound states in the one-dimensional Hubbard model

In this appendix we investigate the nature of the Λ and $k - \Lambda$ strings in the one-dimensional Hubbard model. We show for explicit examples, that both kinds of strings lead to certain kinds of bound states (*i.e.*, the wave function decays exponentially with respect to the differences of coordinates) .

A.1 Λ strings

Let us consider the example of N electrons with spin up and two electrons with spin down forming a Λ string, *i.e.*,

$$\Lambda_2 = \Lambda_1^* \quad , \quad k_1, \dots, k_{N+2} \text{ real} \quad . \quad (\text{A.1})$$

The set $\{\Lambda_1, \Lambda_2 | k_1, \dots, k_{N+2}\}$ must fulfill the periodic boundary conditions (1.11).

The wave function corresponding to this set of spectral parameters is given by

$$\psi_{-1, -1, 1, \dots, 1}(x_1, x_2, \dots, x_{N+2}) = \sum_{P \in S_{2+N}} \text{sgn}(Q) \text{sgn}(P) e^{i \sum_{j=1}^{2+N} k_{P_j} x_{Q_j}} \varphi(y_1, y_2 | P) \quad , \quad (\text{A.2})$$

with amplitudes

$$\begin{aligned} \varphi(y_1, y_2 | P) &= A_{id} F_P(\Lambda_1, y_1) F_P(\Lambda_2, y_2) + A_{21} F_P(\Lambda_2, y_1) F_P(\Lambda_1, y_2) \\ &= A_{id} \left(\prod_{i=1}^{y_1-1} \frac{e_-^{(1)}(P_i)}{e_+^{(1)}(P_i)} \right) \frac{1}{e_+^{(1)}(P_{y_1})} \left(\prod_{i=1}^{y_2-1} \frac{e_-^{(2)}(P_i)}{e_+^{(2)}(P_i)} \right) \frac{1}{e_+^{(2)}(P_{y_2})} \\ &\quad + A_{21} \left(\prod_{i=1}^{y_2-1} \frac{e_-^{(1)}(P_i)}{e_+^{(1)}(P_i)} \right) \frac{1}{e_+^{(1)}(P_{y_2})} \left(\prod_{i=1}^{y_1-1} \frac{e_-^{(2)}(P_i)}{e_+^{(2)}(P_i)} \right) \frac{1}{e_+^{(2)}(P_{y_1})} \quad . \end{aligned} \quad (\text{A.3})$$

We want to show that this wave function decays exponentially with respect to the difference of the coordinates y_1 and y_2 . The only nontrivial y dependent part of

the wave function are the amplitudes φ . Therefore it is sufficient to prove that they decay exponentially.

Taking exponentially small corrections Δ into account, the Λ string is of the form

$$\begin{aligned}\Lambda_1 &= \Lambda + \frac{u + \Delta}{2} \\ \Lambda_2 &= \Lambda - \frac{u + \Delta}{2} \quad \text{with } \Delta^* = -\Delta\end{aligned}\quad . \quad (\text{A.4})$$

As all momenta k_j are real this leads to the following inequalities

$$\left| \frac{e_-^{(1)}(i)}{e_+^{(1)}(i)} \right| > 1 \quad , \quad \left| \frac{e_-^{(2)}(i)}{e_+^{(2)}(i)} \right| < 1 \quad . \quad (\text{A.5})$$

Using the periodic boundary conditions

$$\prod_{i=1}^{N+2} \frac{e_-^{(1)}(i)}{e_+^{(1)}(i)} = \frac{\Lambda_2 - \Lambda_1 - u}{\Lambda_2 - \Lambda_1 + u} = \frac{A_{id}}{A_{21}} \quad (\text{A.6})$$

we can express A_{21} in terms of A_{id} . Using the second set of periodic boundary conditions

$$e^{ik_j L} = \frac{e_-^{(1)}(j)}{e_+^{(1)}(j)} \frac{e_-^{(2)}(j)}{e_+^{(2)}(j)} \quad (\text{A.7})$$

we can express products over $e_{\pm}^{(\alpha)}$ in terms of exponential factors of magnitude 1. Straightforward computations yield

$$\begin{aligned}\varphi(y_1, y_2 | P) &= A_{id} e^{i \sum_{l=1}^{y_1-1} k_{P_l} L} \left(\prod_{i=y_1}^{y_2-1} \frac{e_-^{(2)}(P_i)}{e_+^{(2)}(P_i)} \right) \frac{1}{e_+^{(1)}(P_{y_1})} \frac{1}{e_+^{(2)}(P_{y_2})} + \\ &+ A_{id} e^{i \sum_{l=1}^{y_2-1} k_{P_l} L} \left(\prod_{i=1}^{y_1-1} \frac{e_-^{(2)}(P_i)}{e_+^{(2)}(P_i)} \right) \left(\prod_{i=y_2}^{N+2} \frac{e_-^{(2)}(P_i)}{e_+^{(2)}(P_i)} \right) \frac{1}{e_+^{(1)}(P_{y_2})} \frac{1}{e_+^{(2)}(P_{y_1})} \quad .\end{aligned} \quad (\text{A.8})$$

By definition $y_2 > y_1$ and the inequalities (A.5) ensure that the factor in brackets in the first term (and thus the whole term) in (A.8) decays exponentially for $y_2 \gg y_1$.

The second term can be dropped, because for spin waves N plays the role of the lattice length, and in order to investigate asymptotic properties of the wave function we should consider the infinite volume limit, *i.e.*, $N \longrightarrow \infty$. In this limit the second term can be set to zero as $\left(\prod_{i=y_2}^{N+2} \frac{e_-^{(2)}(P_i)}{e_+^{(2)}(P_i)} \right)$ vanishes.

A.2 $k - \Lambda$ strings

We consider the example $N = M = 2$, *i.e.*, 2 electrons with spin up and 2 with spin down. The periodic boundary conditions read

$$\begin{aligned} e^{ik_j L} &= \frac{e_-^{(1)}(j)}{e_+^{(1)}(j)} \frac{e_-^{(2)}(j)}{e_+^{(2)}(j)} \quad j = 1, \dots, 4 \\ \prod_{j=1}^4 \frac{e_-^{(\beta)}(j)}{e_+^{(\beta)}(j)} &= - \prod_{\alpha=1}^2 \frac{\Lambda_\alpha - \Lambda_\beta - u}{\Lambda_\alpha - \Lambda_\beta + u} \quad \beta = 1, 2 \quad . \end{aligned} \quad (\text{A.9})$$

A $k - \Lambda$ string solution of these equations takes the following form in the $L \longrightarrow \infty$ limit:

$$\begin{aligned} \Lambda_1 &= \Lambda - \frac{u}{2} = \Lambda_2^* \\ k_1 &= \pi - \arcsin(\Lambda - u) \\ k_2 &= \arcsin(\Lambda) = \pi - k_3 \\ k_4 &= \pi - \arcsin(\Lambda + u) \quad . \end{aligned} \quad (\text{A.10})$$

In the finite volume there exist two distinct configurations that both lead to $k - \Lambda$ strings in the limit $L \longrightarrow \infty$, depending on whether k_2 and k_3 are real or complex for finite L .

Case (i): k_2 and k_3 are complex

In this case the k 's can be rearranged such that $\text{Im}(k_1) < \text{Im}(k_2) < 0 < \text{Im}(k_3) < \text{Im}(k_4)$ and the invariance of the periodic boundary conditions under complex conjugation gives the additional constraints

$$k_3^* = k_2 \quad , \quad k_4^* = k_1 \quad . \quad (\text{A.11})$$

Taking this into account one obtains the following solution of (A.9) for finite but large L :

$$\begin{aligned}\Lambda_1 &= \Lambda - \frac{u + \delta}{2} = \Lambda_2^* \\ \sin(k_1) &= \Lambda - u - \frac{\delta}{2} + \epsilon_1 = \sin(k_4^*) \\ \sin(k_2) &= \Lambda + \frac{\delta}{2} + \epsilon_2 = \sin(k_3^*),\end{aligned}\tag{A.12}$$

where δ is purely imaginary and $\epsilon_{1,2}$ are complex. The exponentially small corrections are of the orders

$$\delta = \mathcal{O}\left(2u e^{-i(k_1+k_2^*)L}\right), \quad \epsilon_1 = \mathcal{O}\left(-2u e^{-ik_1L}\right), \quad \epsilon_2 = \mathcal{O}\left(-2u e^{-ik_1L}\right). \tag{A.13}$$

The wave function is given by (A.2) with $N = 2$ and (A.3). After inserting the values of the spectral parameters found in (A.12) into (A.2) and (A.3) and re-normalising the resulting expression one finds :

$$\begin{aligned}\psi_{-1,-1,1,1}(x_1, x_2, x_3, x_4) &= \\ \begin{cases} \operatorname{sgn}(Q)(-1)^{y_1+y_2} \left(e^{i \sum_{j=1}^4 k_j x_{Q_j}} - e^{i \sum_{j=1}^4 k_{R_j} x_{Q_j}} \right) & \text{if } (y_1, y_2) \notin \{(1, 2), (3, 4)\} \\ 0 & \text{else} \end{cases},\end{aligned}\tag{A.14}$$

where R is the permutation $(1, 3, 2, 4)$.

Due to the ordering of the imaginary parts of the momenta k_j both terms on the l.h.s. of (A.14) decay exponentially with respect to the magnitudes of differences of coordinates $|x_k - x_j|$ and thus the wave function describes a bound state.

Case (ii): k_2 and k_3 are real

In this case we must drop the constraint $k_2^* = k_3$. We then find the following solution to (A.9)

$$\Lambda_1 = \Lambda - \frac{u + \delta}{2} = \Lambda_2^*$$

$$\begin{aligned}
\sin(k_1) &= \Lambda - u - \frac{\delta}{2} + \epsilon_1 = \sin(k_4^*) \\
\sin(k_2) &= \Lambda + \epsilon_2 \\
\sin(k_3) &= \Lambda + \epsilon_3 \quad ,
\end{aligned} \tag{A.15}$$

where ϵ_1 is complex and $\epsilon_{2,3}$ are real, while δ is again purely imaginary. The corrections are of the orders

$$\begin{aligned}
\epsilon_1 &= \mathcal{O}\left(-2u e^{-ik_1 L}\right) \\
\delta &= \mathcal{O}\left(\frac{4i \operatorname{Re}(\epsilon_1)}{\cot(\frac{k_2 L}{2}) + \cot(\frac{k_3 L}{2})}\right) \\
\epsilon_2 &= \mathcal{O}\left(\frac{-i}{2} \cot(\frac{k_2 L}{2}) \delta\right) \\
\epsilon_3 &= \mathcal{O}\left(\frac{-i}{2} \cot(\frac{k_3 L}{2}) \delta\right) \quad .
\end{aligned} \tag{A.16}$$

The computation of the wave function is analogous to case (i), the only difference being a new re-normalisation constant. The wave function is given by the same expression as in case (i). Again it describes a bound state although k_2 and k_3 are now real.

B The $M = N = 1$ sector in the Hubbard model

In this appendix we further work out the structure of the Bethe Ansatz wavefunctions in the sector $M = N = 1$. In that sector the wavefunctions depend on parameters k_1 , k_2 and Λ . Our general analysis in sections 3 and 4 gives the following possibilities: (i) we can have $M_e = 2$, $M_1 = 1$, which gives real values for k_1 , k_2 and Λ , or (ii) we can have $M'_1 = 1$, which gives a $m = 1$ k - Λ -string with complex conjugate k_1 , k_2 and real Λ . According to the counting of section 3, we expect to have $\frac{1}{2}L(L-1)$ real solutions (i) and $L-1$ string solutions (ii), giving a total number of $\frac{1}{2}(L-1)(L+2)$ Bethe Ansatz states in this sector. However, we already mentioned that, in the context of the XXX Heisenberg model, there is a ‘redistribution phenomenon’ between different types of solutions, which does not affect the total number of states within a sector,

but which does affect the distribution of those states over various types of string-solutions and real solutions ^[18]. In this appendix we will establish a very similar result for the Hubbard model: we will find that the numbers of the real solutions (i) and string solutions (ii) are not always given by the values quoted above, but that the expected total number $\frac{1}{2}(L-1)(L+2)$ of solutions in this sector can be rigorously established.

Let us consider the Bethe equations for periodic boundary conditions in the sector $M = N = 1$

$$e^{ik_1 L} = \frac{\sin k_1 - \Lambda - \frac{u}{2}}{\sin k_1 - \Lambda + \frac{u}{2}}, \quad e^{ik_2 L} = \frac{\sin k_2 - \Lambda - \frac{u}{2}}{\sin k_2 - \Lambda + \frac{u}{2}}$$

$$\prod_{i=1}^2 \frac{\sin k_i - \Lambda - \frac{u}{2}}{\sin k_i - \Lambda + \frac{u}{2}} = 1. \quad (\text{B.1})$$

Since we assume Λ to be finite, we can solve for it and find $\Lambda = \frac{1}{2}(\sin k_1 + \sin k_2)$. The equations then reduce to

$$e^{ik_1 L} = \frac{\sin k_1 - \sin k_2 - u}{\sin k_1 - \sin k_2 + u}, \quad e^{i(k_1+k_2)L} = 1. \quad (\text{B.2})$$

We can solve the second equation by putting $k_1+k_2 = \frac{2\pi}{L}m$, with $m = 0, 1, \dots, 2L-1$ and write $k_1 = \frac{\pi}{L}m + x$ and $k_2 = \frac{\pi}{L}m - x$. The remaining equation reads

$$e^{i(\pi m + Lx)} = \frac{-\frac{4}{U} \cos(\frac{\pi m}{L}) \sin x - i}{-\frac{4}{U} \cos(\frac{\pi m}{L}) \sin x + i}. \quad (\text{B.3})$$

One easily checks that, if x_0 solves this equation for $m = m_0$, then $x = x_0 + \pi$ solves the equation for $m = m_0 + L$, and that the resulting wavefunctions are the same. We can thus restrict our attention to $m = 0, 1, \dots, L-1$ and $-\pi \leq x < \pi$.

Let us now try to find real solutions x for the equation (B.3) for given m . Taking a logarithm we have

$$\arctan\left(-\frac{U}{4} \cos(\frac{\pi m}{L}) \sin x\right) = \frac{1}{2}Lx - \pi n, \quad (\text{B.4})$$

where n is an arbitrary integer for m odd and half an odd integer for m even. It is rather straightforward to solve this equations by a graphical method: one plots both

the l.h.s and the r.h.s. (for various n) of these equations on the interval $-\pi \leq x < \pi$ and reads of the intersection points, which are then solutions of the equation. For large enough $|U|$ this procedure is easily carried out and one finds the following.

For m even there are solutions for $n = -\frac{L+1}{2}, \dots, \frac{L-1}{2}$, which are L solutions in total. Since solutions x and $-x$ are equivalent (and $x = 0$ is not among the solutions) we should divide this number by 2. Using that there are $L/2$ possible even values for m we thus find $L^2/4$ solutions. For m odd one finds non-equivalent solutions for $n = 1, 2, \dots, \frac{L-1}{2}$ for each m , which gives a total number of $(L^2 - 2L)/4$. (The solutions $x = 0$, $x = \pm\pi$, which exist for generic U , give vanishing wavefunctions in general.) Adding up the contributions from odd and even m , we find $\frac{1}{2}L(L-1)$, which is indeed the number predicted by the counting in section 3.

However, let us now assume that m is odd and that U is close to a critical value U_m , which we define by

$$\frac{U_m}{4} \cos\left(\frac{\pi m}{L}\right) = \frac{L}{2}. \quad (\text{B.5})$$

At the value $U = U_m$ the curve for the l.h.s. of (B.4) has slope $L/2$ at $x = \pm\pi$, and at the value $U = -U_m$ the curve for the l.h.s. of (B.4) has slope $L/2$ at $x = 0$. Since the r.h.s. is given by straight lines of slope $L/2$, and since both curves already had intersections at $x = 0$ (for $n = 0$) and $x = \pm\pi$ (for $n = \pm L/2$) (which did however not give rise to non-trivial wavefunctions), it will be clear that the number of intersections changes when U reaches the critical values $\pm U_m$. In fact, one finds one extra real solution x (together with the equivalent solution $-x$) for a given odd m as soon as $U < |U_m|$. For example, if U is such that $U_1 > U > U_3 > \dots > 0$ there will be one extra real solution to the equations (B.2).

The complex values for x which solve the equation (B.3) are of the form $x = iy$ or $x = \pi + iy$ with y real. In a way similar to what we showed above, one can analyse the equations for the real quantity y by a graphical method. If $|U|$ is sufficiently large, one finds precisely one complex solution for $m = 1, 2, \dots, L$ with the exception of $m = L/2$. In that case, there are thus $L-1$ complex solutions, which is in agreement with the counting of section 3.

However, from the graphical analysis one finds that the complex solution for a given odd m disappears as soon as $|U|$ is chosen to be smaller than U_m . Note that

this happens precisely in the regime where we have found one extra real solution!

We thus find a redistribution phenomenon, where solutions change their nature as a function of U , in analogy to what we found for the XXX Heisenberg model in [18]. In the Hubbard model the phenomenon is easily understood: if $|U|$ is made small enough, the interactions become so weak that some of the bound states (with complex x) decay into real solutions (with real x).

When $|U|$ is chosen to be equal to one of the critical values U_m , there do exist nontrivial wavefunctions with $x = 0$ or $x = \pm\pi$, *i.e.*, with coinciding k_1 and k_2 . These wavefunctions can be seen to be nonvanishing by a renormalisation à la l'Hôpital.

In all cases, the total number of eigenfunctions of the hamiltonian in the sector $M = N = 1$ is found to be $\frac{1}{2}(L-1)(L+2)$, which is the value predicted by the counting in section 2, and used for the proof of completeness in section 3.